On the Area Bisectors of a Polygon*

Karl-Friedrich Böhringer† University of California, Berkeley

Bruce Randall Donald‡ Dartmouth College

Dan Halperin§ Tel Aviv University

Abstract

We consider the family of lines that are area bisectors of a polygon (possibly with holes) in the plane. We say that two bisectors of a polygon P are *combinatorially distinct* if they induce different partitionings of the vertices of P. We derive an algebraic characterization of area bisectors. We then show that there are simple polygons with n vertices that have $\Omega(n^2)$ combinatorially distinct area bisectors (matching the obvious upper bound), and present an output-sensitive algorithm for computing an explicit representation of all the bisectors of a given polygon. Our study is motivated by the development of novel, flexible feeding devices for parts positioning and orienting. The question of determining all the bisectors of polygonal parts arises in connection with the development of efficient part positioning strategies when using these devices.

*Work on this paper by Karl-Friedrich Böhringer and Bruce Randall Donald has been supported in part by the National Science Foundation under grants No. IRI-8802390, IRI-9000532, IRI-9201699, and by a Presidential Young Investigator award to Bruce Donald, in part by NSF/ARPA Special Grant for Experimental Research No. IRI-9403903, and in part by the Air Force Office of Sponsored Research, the Mathematical Sciences Institute, Intel Corporation, and AT&T Bell laboratories. Work on this paper by Dan Halperin has been supported in part by an Alon Fellowship, by ESPRIT IV LTR Project No. 21957 (CGAL), by the USA-Israel Binational Science Foundation, and by the Hermann Minkowski – Minerva Center for Geometry at Tel Aviv University. A preliminary and abridged version of the paper appeared in *proc. 13th ACM Symp. on Computational Geometry*, Nice, 1997, pp. 457–459.

†ALPHA laboratory, Dept. of Ind. Eng. and Op. Research, University of California, Berkeley, CA 94720-1777, karl@IEOR.Berkeley.EDU, http://www.ieor.berkeley.edu/~karl.

‡Dept. of Computer Science, Dartmouth College, 6211 Sudikoff Laboratory, Hanover, NH 03755-3510. brd@cs.dartmouth.edu, http://www.cs.dartmouth.edu/~brd.

§Department of Computer Science, Tel Aviv University, Tel Aviv 69978, ISRAEL, halperin@math.tau.ac.il. Part of the work on this paper was carried out while D.H. was at the Robotics Laboratory, Department of Computer Science, Stanford University.

1 Introduction

Let P be a polygon in the plane, possibly with holes, and having n vertices in total. We denote by V the set of vertices of P. For a directed line λ in the plane, we denote by $h_l(\lambda)$ (resp. $h_r(\lambda)$) the open half-plane bounded by λ on the left- (resp. right-) hand-side of λ. The line λ is an area bisector of P if the area of $P \cap h_l(\lambda)$ is equal to the area of $P \cap h_r(\lambda)$.

A line λ partitions V into three sets (two of which may be empty): $V \cap h_l(\lambda)$, $V \cap \lambda$, and $V \cap h_r(\lambda)$. We say that two area bisectors of P are *combinatorially distinct* if the partitioning of V as above induced by the two bisectors is different. We say that two area bisectors of P are *combinatorially equivalent* if they induce the same partitioning of V.

We assume that the polygon P is connected, and non-degenerate in the sense that its interior is connected.

If instead of a polygon we take a set of n points, we can ask how many combinatorially distinct *halving lines* there are, namely, instead of equating the area on the two sides of a line we now wish to equate the cardinality of the subsets on either side of the line. This is the well-known *k-set* problem (for $k = \frac{n}{2}$) that has been extensively studied in discrete geometry [8]. The best bounds known to-date for this problem are the recently obtained upper bound $O(n^{4/3})$ [5] and a lower bound $\Omega(n \log n)$ [10]. An obvious upper bound on the number of distinct *area* bisectors of a polygon with n vertices is $O(n^2)$—see Section 2. In this paper we show that a polygon with n vertices can have $\Omega(n^2)$ distinct area bisectors. (Note that the polygon in our construction is *simple*.)

We devise an output-sensitive algorithm for computing an explicit representation of all the area bisectors of a given polygon. In our analysis we give a more refined distinction between area bisectors (than the distinction by the partitioning of V). We order the area bisectors by slope in the range $[-\pi/2, \pi/2)$, and partition the range into maximal connected intervals with bisectors of the same combinatorial equivalence class. We denote by $\kappa = \kappa(P)$ the number of such slope intervals for a polygon P, and by $\kappa(n)$ the maximum number of such slope intervals over all non-degenerate polygons with n vertices. We denote by $K = K(P)$ the number of combinatorially distinct area bisectors of a polygon P, and by $K(n)$ the maximum number of combinatorially distinct area bisectors over all non-degenerate polygons with n vertices. It is clear that $\kappa(n) \geq K(n)$; we show below that there are polygons for which $\kappa(P) > K(P)$.

It is convenient to consider the problem at a dual setting as well. The *bisector curve* β is defined in a plane dual to the plane containing the polygon: β is the union of points dual to area bisectors. We show that β is x-monotone in the dual plane (where the x-coordinate corresponds to the slope of the bisector) and that it is a continuous piece-wise algebraic curve. Each maximal piece b of β that is defined by the same function describes a contiguous (in slope) set of bisectors that cross the same set of edges of the polygon P, the endpoints of b correspond to bisectors that contain a vertex of P, and any bisector described by an interior point of b does not contain any vertex of P. Hence, $\kappa(P)$ denotes the number of maximal algebraic pieces of β for a polygon P.

A c-oriented polygon is a polygon all whose edges are parallel to exactly c directions, where c is a fixed constant. For a c-oriented polygon with n vertices, our algorithm for computing an explicit representation of the bisectors by determining the pieces of the curve β runs in time $O((n+\kappa)\log^2 n)$. For a general non-degenerate polygon having n vertices the

algorithm runs in time $O((n+\kappa)\log^2 n+(n+\kappa)\alpha(n)\psi(t))$. Here and throughout the paper, $\alpha(\cdot)$ denotes the extremely slowly growing functional inverse of Ackermann's function, t denotes the maximum number of distinct slopes of edges of P intersected by any line, and $\psi(j)$ denotes the time required to find the roots of a polynomial equation of degree j.

Our algorithm proceeds by constructing the *zone* of the curve β in an *arrangement of lines* in a plane dual to the plane of the polygon. An arrangement $\mathcal{A}(L)$ of a set L of lines is the subdivision of the plane induced by L into vertices, edges, and faces [8]. The zone of a curve in an arrangement of lines is the collection of faces of the arrangement crossed by the curve [12]. In our case the equations defining the curve β depend on the face of the arrangement that β crosses, that is, these equations change from face to face. Hence, we could not use ready-made algorithms for computing the zone of β since the algorithms we are aware of assume that the curve for which the zone is computed is known in advance.

Area bisectors were considered by Díaz and O'Rourke [6, 7]. However, their focus is on the continuous version of the *Ham-sandwich cut* problem, and of a problem they introduce of *orthogonal four-sections*; see [6, 7] for more details.

Our study is motivated by the development of novel, flexible feeding devices for parts positioning and orienting. These devices are based on the principle of *programmable force fields* [3]. They often exploit exotic actuation technologies such as MEMS (micro electro mechanical system) actuator arrays [4] or transversely vibrating plates [2] to generate controllable planar fields of 2d force vectors. When a part is placed on these devices, the programmed vector field induces a force and moment upon it. Over time, the part may come to rest in a dynamic equilibrium state. By chaining together sequences of vector fields, the equilibria may be cascaded to obtain a desired final state—for example, this state may represent a unique orientation or pose of the part. Figure 1 shows a strategy for manipulating a part to a unique orientation equilibrium (up to symmetry), using a sequence of *squeeze fields.*

A squeeze field is defined as a force vector field that, at each point, has a unit force vector pointing perpendicularly towards a common *squeeze line*. Such a field can be implemented for example by a MEMS actuator array, by making all the actuators push in the appropriate direction. In order to be in equilibrium, the forces and moments acting on a part must balance. Force equilibrium, under reasonable assumptions detailed in [3], is equivalent to the following: *the squeeze line must be an area bisector of the polygon.* Similarly, moment equilibrium requires that the centers of area of the left and right part sections lie on a line perpendicular to the bisector. Currently, our algorithm for equilibrium analysis requires the explicit identification of all the area bisectors of a part. The resulting manipulation plans depend on the number of distinct bisectors, motivating the problems discussed in this paper. For more details see [1, 3, 4].

The rest of the paper is organized as follows. In Section 2 we present basic properties of area bisectors that are later necessary for the development of the algorithm. We postpone most of the technical analysis in this section to the Appendix. In Section 3 we present a simple polygon with n vertices that has $\Omega(n^2)$ combinatorially distinct area bisectors. The algorithmic results are then presented in Section 4. In Section 5 we give concluding remarks and point to open problems raised by our study.



Figure 1: Sensorless part alignment: The part reaches unique orientation after two subsequent squeezes. There exist such alignment plans for all polygonal parts. See URL `http://www.cs.cornell.edu/home/karl/MicroManipulation` for an animated simulation. Figure reprinted from [4].

2 Properties of Area Bisectors

In this section we show that all combinatorially equivalent bisectors of a fixed equivalence class of any polygon can be described by a rational function. We develop explicit formulas for bisectors for given polygon geometry and investigate their algebraic complexity—this is summarized in Propositions 2 and 3 below. For ease of exposition we postpone most of the technical algebraic analysis to the Appendix.

We extend the definition of *combinatorial equivalence* (see Introduction) to arbitrary lines that partition a polygon. We define combinatorially equivalent partitioning lines of a polygon P to be all those lines λ that induce the same vertex partition $V \cap h_l(\lambda)$, $V \cap \lambda$, and $V \cap h_r(\lambda)$. A necessary (but not sufficient) condition for combinatorial equivalence is that λ intersects the same ordered set of polygon edges.

It is convenient to study the problem in a dual plane: a line $y = 2\bar{x}x - \bar{y}$ in the primal plane is transformed into the point $(\bar{x}, \bar{y})$ in the dual plane. A point (x, y) in the primal plane is transformed into the line $\bar{y} = 2x\bar{x} - y$ in the dual. The dual of an object o will be denoted by o^*. If O is a set of objects in the plane, O^* will denote the set of dual objects. For details on duality transforms see, e.g., [14].

In the following lemma we summarize basic facts on area bisectors of a polygon.

Lemma 1 *Given a polygon P with vertices V, $n := |V|$, and a line λ.*

1. *There exist $O(n^2)$ combinatorially distinct ways in which a line can partition P.*
2. *Let A and B be the intersections of a bisector λ with the boundary of the convex hull of P. As the slope $\bar{x}$ of λ grows from $-\pi/2$ to $\pi/2$, A and B progress monotonically counterclockwise on the boundary of the convex hull of P.*
3. *If the interior of P is connected, then there exists a unique bisector of P for every*



Figure 2: Two non-parallel lines λ and λ' in combinatorially equivalent intersection with polygon P.

slope $\bar{x}$ of λ.

Proof: 1. This is a well-known fact. We prove it here to demonstrate how duality is used in our arguments. The dual V^* of the n polygon vertices V form an arrangement of n lines in the dual plane. Each point on a line v^* in the dual plane corresponds to a line through the polygon vertex v in the primal plane, and vice versa. Each point within a face of the dual arrangement corresponds to a line in the primal plane that does not intersect any polygon vertices. Thus, all points within a particular face of the dual arrangement correspond to combinatorially equivalent line placements. Since an arrangement of n lines has $O(n^2)$ faces, there are at most $O(n^2)$ combinatorially distinct ways in which a line can partition P.

Proofs of items 2 and 3 can be found in Díaz's thesis [7]. □

Now consider a bisector λ of polygon P for varying $\bar{x}$ values, as described in Lemma 1.2. The intersections of λ with the boundary of the convex hull of P, A and B, progress monotonically about that boundary. In general, this progression corresponds to a simultaneous rotation and translation of λ.

Our goal is to obtain a precise mathematical relationship between bisector slope $\bar{x}$ and intercept $\bar{y}$. To this end, we first analyze the effects of rotation and translation separately. To simplify the analysis, we choose a parameterization of the partitioning line λ different from $\bar{x}$ and $\bar{y}$. The line λ is given by a fixed point s on λ and a vector p that specifies its direction (see Figures 2 and 3).[1]

The detailed analysis is given in the Appendix and its results are given in Proposition 2 below. Note that during pure rotation or pure translation, in general the bisector property is not maintained. By imposing the bisector condition $|P \cap h_l(\lambda)| = |P \cap h_r(\lambda)|$ we obtain the

[1] The transformation to $\bar{x}$ and $\bar{y}$ is straightforward: Given $s = (s_x, s_y)$ and $p = (p_x, p_y)$, we get $\bar{x} = \frac{p_y}{2p_x}$ and $\bar{y} = \frac{p_y s_x}{p_x} - s_y$.



Figure 3: Two parallel lines λ' and λ'' in combinatorially equivalent intersection with polygon P.

relationship between rotation and translation of λ. This will be addressed in Proposition 3.

Proposition 2 *Let P be a polygon with n vertices. Let s be a point in $\mathbb{R}^2$ and p be a vector in $\mathbb{R}^2$, and let λ be a line through s in direction p that intersects r edges of P with $t \leq r$ distinct slopes.*

1. *Let λ' be a line through s in direction p', where $p' = p + \alpha q$ for some fixed vector q and some parameter $\alpha \in \mathbb{R}$. If λ' is combinatorially equivalent to λ, then the area of $P \cap h_l(\lambda')$ is given by a rational function $a_l(\alpha)$ of numerator and denominator degree $O(t)$. In particular, the combinatorially equivalent area bisectors of P passing through s are determined by the roots of a polynomial equation of degree t.*
2. *Let λ'' be a line parallel to λ' with distance α' to λ'. If λ'' is combinatorially equivalent to λ', then the area of $P \cap h_l(\lambda'')$ is given by a quadratic polynomial $b_l(\alpha')$.*

A detailed proof and complete formulas for Proposition 2 can be found in the Appendix. Note that the rational functions a_l and b_l express the changing area of $P \cap h_l(\lambda)$ for pure rotation and pure translation of λ, respectively. If λ is a bisector, this area needs to remain constant, which results in a coupling between the rotational parameter α and the translational parameter α'. This relationship is characterized in the following Proposition:

Proposition 3 *Let λ be a bisector of P, and consider the combinatorially equivalent line λ'' parameterized by $\alpha, \alpha' \in \mathbb{R}$ as described in Proposition 2 above. There exists a function $f(\alpha, \alpha')$ such that λ'' is a bisector of P iff $f(\alpha, \alpha') = 0$. f is a polynomial of degree t in α and a quadratic polynomial in α', where $t \leq r$ is the number of distinct slopes of the r polygon edges intersected by λ and λ''.*

The polynomial $f(\alpha, \alpha')$ is given in the Appendix (Equation 8), which includes a detailed derivation of this result. With Proposition 3 we obtain a parametric description for every set



Figure 4: A simple polygon with n vertices for which $\kappa > K$: By continuously changing the slope of λ, a particular equivalence class of bisectors is entered and exited $\Omega(n)$ times.

of combinatorially equivalent bisectors of P: Given a polygon, we can generate Equation 8 for any particular equivalence class, which yields the polynomial equation $f(\alpha, \alpha') = 0$. Solving f for α', we finally obtain $\bar{x}$ and $\bar{y}$ parametrized by α.

While Proposition 3 gives an algebraic characterization of the area bisectors, in subsequent sections we will only be using an intermediate result, summarized in Proposition 2.1, describing the area bisectors through a given point in the plane. In this case we have a univariate (in α) polynomial equation. We will use this equation in the time analysis of the algorithm in Section 4, and for a combinatorial upper bound on κ in Section 5.

We conclude this section by showing that there are polygons P for which $\kappa(P)$ is strictly greater than $K(P)$. This example is due to Lovett.[2] We show that for a given polygon with n vertices and two combinatorially equivalent area bisectors λ_1 and λ_2, as we vary the slope between the slope of λ_1 and λ_2 the corresponding area bisectors exit and enter the equivalence class of λ_1 (and λ_2) $\Omega(n)$ times.

Let Δabc be a triangle. It is easily verified that a median in a triangle is an area bisector of the triangle. We denote the median incident to c by λ_1, and the median incident to a by λ_2. Consider the family F of all the area bisectors obtained when we vary the slope between the slopes of λ_1 and λ_2, in the interval where all area bisectors intersect the edges ab and bc. We take the upper envelope of the lines in F, which is a convex curve E.[3] See Figure 4 for an illustration. For a given n, we take a set Q of $\lfloor (n-1)/3 \rfloor$ evenly spaced points along E, and draw the tangent line to E at each of these points. Each tangent line

[2] S. Lovett, personal communication, Tel Aviv University, 1997.

[3] For example, if we choose the coordinates of T's vertices to be $a = (0, 1), b = (0, 0)$, and $c = (2, 0)$, then E is the hyperbola $f(x) = \frac{1}{4x}$.



Figure 5: A simple polygon with n vertices that has $\Omega(n^2)$ combinatorially distinct bisectors.

is an area bisector. Consider two consecutive points and their tangents: In between the points, slightly below the envelope and above the two tangent lines we draw the apex of a thin spike extending from the edge ac of the triangle. We carve this spike out of the triangle. We repeat this process for each pair of consecutive points of $Q \cup \{a, c\}$. Finally we pull the edge ac outwards so that the area of the polygon above λ_1 (which is half the area of the original triangle) will be equal to the area below λ_1, compensating for the area of the spikes.

All the tangent lines induced by the set Q are area bisectors of the same equivalence class—they have the single vertex b on one side of the bisector. Evidently as we let the slope vary, every time we cross a spike the equivalence class changes. Hence we obtained a polygon with n vertices and an equivalence class of area bisectors that is entered and exited $\Omega(n)$ times as we let the slope vary.

3 Lower Bound

As argued above, a polygon with n vertices can have at most $O(n^2)$ combinatorially distinct area bisectors. In this section we give an example of a simple polygon with n vertices where the bound $\Omega(n^2)$ is attained.

Consider Figure 5. All the vertices v_i, v'_i, u_i and u'_i lie on a circle whose center is at c. The vertices w_j lie very close to c on a small circle whose center is c as well, along two convex polygonal chains.

We fix an integer m (that we will determine later; for the polygon in the figure $m = 3$).

The distance between the vertices v_i and v_{i+1} is the same for $i = 1, \ldots, m$, and it is the same as the distance between v'_i and v'_{i+1} for $i = 1, \ldots, m$. The area of all triangles $v_i u_i v_{i+1}$ for $i = 1, \ldots, m$ is the same and is equal to the area of all triangles $v'_i u'_i v'_{i+1}$ for $i = 1, \ldots, m$. There are $2m$ vertices w_j near c and they are equally spaced on a small circle centered at c. As can be easily verified, for every pair of vertices v_i and v'_i, there is a bisector passing through these points that passes also through the center point c. We next claim that as we rotate the bisector from v_i to v_{i+1} it will move off the center c and sweep m vertices w_j.

The reason that the bisector will move off center as it rotates from v_i to v_{i+1} is that the angle $\angle u_i v_i v_{i+1}$ is greater than the angle $\angle u'_i v'_i v'_{i+1}$. Hence, as the bisector rotates, it will proceed 'faster' on the bottom part of our polygon than on the top part and therefore will sweep half of the vertices w_j on its way.

Finally m is chosen such that (roughly) $n = 6m + 8$. It follows that the number of distinct area bisectors is $\Omega(m^2) = \Omega(n^2)$.

4 Output-Sensitive Algorithm

Let P be a polygon with n vertices as defined in the Introduction, namely connected, non-degenerate and possibly with holes. Throughout this section we assume that no three vertices of P are collinear (see also remark at the end of the section). In the dual plane every vertex v of P is transformed into a line v^* which is the collection of all points dual to lines in the primal plane that pass through v.

For any given direction there is a unique area bisector. We denote the oriented bisector of P that makes an angle θ with the positive x-axis by $B(\theta)$, and (because of symmetry) confine ourselves to the range $[-\pi/2, \pi/2)$ for θ. We denote the collection of points dual to area bisectors of P in that range by β. Note that any θ (besides $-\pi/2$) corresponds to an $\bar{x}$-coordinate in the dual plane.

The curve β is a piece-wise algebraic and $\bar{x}$-monotone curve (see Proposition 3 and Lemma 1.3). We call β the *bisector curve* of P, as it gives a complete specification of all the area bisectors of the polygon P. The number κ as defined in the Introduction corresponds to the number of maximal connected algebraic pieces in β, where the function describing each piece is defined by the fixed set of edges that the corresponding set of bisectors cross. In this section we describe an output-sensitive algorithm to compute β.

Since we aim for output-sensitivity, we cannot afford to compute the entire arrangement $\mathcal{A}(V^*)$ whose complexity can be $\Omega(n^2)$. We will discover the maximal pieces of β in their order along β by dynamically exploring the faces of the arrangement through which β passes, and by computing the intersection points of β with the edges bounding these faces.

We choose an arbitrary direction $\theta_0 \in [-\pi/2, \pi/2)$ and look for the area bisector of P in that direction. This can be done in $O(n \log n)$ time for an arbitrary polygon [6], and in $O(n)$ time for a simple polygon [16]. We assume that the bisector $B(\theta_0)$ does not pass through a vertex of P, otherwise we change θ_0 slightly so that $B(\theta_0)$ does not pass through a vertex.

Next, we obtain the set of edges crossed by $B(\theta_0)$ in $O(n)$ time (this could also be obtained as a by-product of the algorithm for finding $B(\theta_0)$). We denote by $E(\theta)$ the set



Figure 6: (a) The face f containing p is the intersection of half-planes containing p; (b) the maximal pieces of β inside f and its exit points from f.

of edges crossed by $B(\theta)$. The set $E(\theta_0)$ determines the function describing the bisector curve β in a neighborhood of θ_0. More precisely, the set $E(\theta_0)$ induces a function that describes the bisectors as long as the set of edges crossed by the bisector does not change (see Section 2). In the dual plane this function describes the curve β as long as we do not leave the face of $\mathcal{A}(V^*)$ which contains the point $p := (B(\theta_0))^*$.

Our next step is to construct the face $f = f(p)$ that contains the point p in $\mathcal{A}(V^*)$. We will do a little more than just exploring the face f, in preparation for the rest of the algorithm. For each line $v_i^* \in V^*$, let $h_i(f)$ denote the half-plane bounded by v_i^* and containing the face f. Let $H = H(f)$ denote the collection of n half-planes $h_i(f)$. The face f equals the intersection of all half-planes in $H(f)$; see Figure 6 for an illustration. We will explore the faces of the zone of β in the arrangement $\mathcal{A}(V^*)$ by dynamically maintaining the intersection of half-planes H [15]. We denote the necessary dynamic data structure by D. The time to construct the face f and prepare D is $O(n \log^2 n)$ [15].

Now we determine the maximal pieces of $\beta \cap f$. We split the edges bounding f into an upper chain and a lower chain, each set being ordered from left to right. For each edge on the upper chain we compute the intersection of its supporting line with β. Computing the intersection of β with the line v^* supporting an edge is equivalent to finding the bisectors that pass through the vertex v and intersect a fixed set of edges of P. Hence we can use Equation 4 (see also Proposition 2.1). The roots of the polynomial determine the intersection points; we may have to filter out roots that correspond to bisectors with slope outside the given slope interval. We order the resulting intersections along the $\bar{x}$-axis. We repeat the same for the edges of the lower chain, and merge the two lists of intersection points into a single list I. Since the curve β is $\bar{x}$-monotone, the list I provides a description of the curve β inside f. Moreover, the list I indicates what are the neighboring faces that β crosses. We mark each of these additional faces by the point where β crosses out of f. We call each such point an *exit point*. See Figure 6(b).

Since f has already been constructed, we know for each exit point of β the line that contains it (in fact we know a little more, we also know the two lines that bound the edge from which β exits). We next explore one of the faces, say f', into which the curve β crosses out of f through an edge that we denote by e. Let v_i^* be the line containing the edge e. In terms of intersection of half-planes, to obtain f' the only half-plane that needs to be modified is the half-plane supported by v_i^*. More specifically to obtain the new set $H(f')$,

we delete $h_i(f)$ from H and add $h_i(f')$ which is the half-plane supported by v_i^* on the other side of v_i^*. (Some caution is needed if β crosses out of f through a vertex. However, since only two lines intersect at that vertex, the operations are similar to those described for the case of crossing out of an edge, and their asymptotic running time is the same.)

Thus we need to delete a half-plane from D and insert a half-plane into D. As a result we obtain the new face f'. The cost of the operation is $O(\log^2 n + C(f'))$, where $C(f')$ is the number of edges on the boundary of f'. The insertion and deletion each costs $O(\log^2 n)$, and the cost of reporting the edges on the boundary of the new face f' is proportional to the number of these edges [15]. We keep a data structure, say a quad-edge structure [11], that describes all the faces of $\mathcal{A}(V^*)$ that have already been constructed so that we do not construct the same face twice. The cost of updating the quad-edge structure with the new face is proportional to $C(f')$.

We continue exploring the faces through which β passes by moving monotonically in the positive x-direction. When the move in that direction has been completed, namely there are no more exit points to the right of the last exit point handled, we return to the starting point p and repeat the process in the negative x-direction. The algorithm stops when we have identified all the intersection points of β with lines in V^*, and so we have also identified the *zone* of β in $\mathcal{A}(V^*)$, namely all the faces of $\mathcal{A}(V^*)$ crossed by β.

Recall that κ is the sum, over all faces f in the zone of β in $\mathcal{A}(V^*)$, of the number of maximal connected components $\beta \cap f$. The overall running time of the algorithm consists of the following components.

1. The initial construction of the structure D takes $O(n \log^2 n)$.
2. For each exit point we pay $O(\log^2 n)$ to update D, for a total of $O(\kappa \log^2 n)$.
3. For each newly visited face f we also pay $O(C(f))$ to report its bounding edges and keep them in the quad-edge structure.
4. For each edge on each explored face we compute its intersection with β.

To estimate the cost of components 3 and 4, we need a bound on the sum of $C(f)$ over all faces f of the zone of β in the arrangement $\mathcal{A}(V^*)$. The following lemma bounds the complexity of the zone of an arbitrary curve in an arrangement of lines. We make no assumptions on the curve besides that it consists of a single connected component.

Lemma 4 *Given a collection $\mathcal{L}$ of n lines in the plane and a connected curve γ that intersects the lines of $\mathcal{L}$ in N points in total, then the complexity of the zone of γ in $\mathcal{A}(\mathcal{L})$ is $O((n+N)\alpha(n))$.*

Proof: We modify the arrangements $\mathcal{A}(\mathcal{L})$ as follows (by an idea borrowed from [9]). Wherever γ crosses a line $\ell \in \mathcal{L}$, we make a small gap in ℓ, splitting ℓ into two segments such that γ passes between them. As a result we get a collection S of $n+N$ segments, and γ lies in a single face of the arrangement $\mathcal{A}(S)$. Since the complexity of a single face in an arrangement of t segments is $O(t\alpha(t))$ a bound of $O((n+N)\alpha(n+N))$ follows. Har-Peled [13] showed that if all the segments lie on only n lines then the complexity of a single face is $O((n+N)\alpha(n))$. □

Since the number of intersection points between β and V^* is $\kappa - 1$ we have the following immediate corollary.

Corollary 5 *The maximum combinatorial complexity of the zone of β in $\mathcal{A}(V^*)$ is $O((n+\kappa)\alpha(n))$.*

Thus component 3 of the running time is $O((n+\kappa)\alpha(n))$ which is subsumed by components 1 and 2. It remains to bound component 4, namely, the time to compute all the intersection points of β with the edges of the zone of β. Let t denote the maximum number of distinct slopes of edges of P intersected by a single line. Also, let $\psi(j)$ denote the time to find the roots of a polynomial equation of degree j. For general polygons (of the type defined in the Introduction) the time to compute all the intersection points of β with V^* is obviously bounded by $O((n+\kappa)\alpha(n)\psi(t))$.

If P is c-oriented, then by Proposition 2.1 the equation describing the bisectors through a given vertex and intersecting a fixed set of edges is a polynomial of constant maximum degree and it takes constant time to find its roots. It follows that, in this case, the time to compute the intersections of β with the lines V^* is asymptotically dominated by the time to compute the zone of β in $\mathcal{A}(V^*)$.

We summarize the discussion above in the following theorem.

Theorem 6 *Given a non-degenerate polygon P (possibly with holes) with n vertices, then we can find a complete specification of its bisectors in time $O((n+\kappa)\log^2 n + (n+\kappa)\alpha(n)\psi(t))$, where κ is the number of maximal connected algebraic pieces of the bisector curve β. If P is c-oriented, then the algorithms runs in time $O((n+\kappa)\log^2 n)$.*

Remark. We assumed above that no three vertices of the polygon P are collinear. If we relax this assumption the following situation may arise. As we explore the zone of β in the arrangement, β crosses out of a face f through a vertex u at which $n_u > 2$ lines are incident. We could reach the face into which β crosses out of f by $O(n_u)$ update operations to the structure D. However, the overall cost of operations at such degenerate crossings is no longer as sharply related to κ as in the bounds above.

5 Conclusions and Open Problems

In this paper we have characterized the family of area bisectors of a polygon. We have shown that a polygon with n vertices can have $\Theta(n^2)$ combinatorially distinct area bisectors, namely $\Theta(n^2)$ equivalence classes of bisectors, where two bisectors are equivalent if the lines containing them split the set of polygon vertices in the same fashion. We have also presented an output-sensitive algorithm that produces an explicit representation of all bisectors of a polygon P.

Our algorithm produces the bisector families ordered by slope. We have shown at the end of Section 2 that there exist polygons for which, as we vary the slope of the bisector, the same equivalence class can be entered and exited $\Omega(n)$ times. This raises the following problem: How many equivalence classes of bisectors can be encountered when we let the slope of the bisector vary in the range $[-\pi/2, \pi/2)$; in the paper's notation the question is to obtain a bound on the value $\kappa(n)$. The lower bound $\Omega(n^2)$ for $K(n)$ shown in section 3

applies here as well. The best upper bound we can show at the moment is $O(n^3)$ derived by the following argument, using the dual arrangement formulation. Consider one edge e of the arrangement $\mathcal{A}(V^*)$. The bisectors whose dual points lie on e are determined by the roots of a polynomial of degree at most n (Equation 4), so their number is at most n, thus the bisector curve cannot cross an edge of the arrangement more than n times. Since the arrangement has $O(n^2)$ edges, the cubic bound follows. In summary there is a gap of an order of magnitude between the lower bound and upper bound on $\kappa(n)$ and the problem is to tighten this gap.

Another open problem is raised by the motivating application. As explained in the Introduction, area bisectors correspond to force equilibria of polygonal parts put on certain part-orienting devices. We are in fact interested in *total* equilibria which are simultaneously force and *moment* equilibria of polygonal parts. A line λ crossing a polygon P induces total equilibrium if (i) λ is an area bisector of P, and (ii) the line connecting the center of mass (area) of $P \cap h_l(\lambda)$ with the center of area of $P \cap h_r(\lambda)$ is perpendicular to λ. The problems we propose are to bound the number of total equilibria of a polygon, and to efficiently compute them. For the combinatorial bound we have only trivial bounds: a lower bound $\Omega(n)$ for a regular n-gon and an upper bound $O(n^3)$ which is obtained in a way similar to the upper bound for $\kappa(n)$, namely for every equivalence class of area bisectors, the total equilibria are defined by the roots of a polynomial of degree at most n, where n is the number of vertices of the polygon. As for an algorithm, our way to obtain total equilibria is first to find all the distinct bisectors and then solve the problem for each class of bisectors. Could the stage of computing all bisectors be avoided when computing total equilibria? For example, for a sufficiently large n, the part in Figure 5 (Section 3) has only 1 stable (and 1 unstable) total equilibrium, even though there exist $\Omega(n^2)$ distinct area bisectors.[4] This equilibrium is attained when the connecting bar between the outer circle segments and the inner circle is parallel to the squeeze line, and for the unstable equilibrium, the squeeze line is perpendicular. In both cases the squeeze line passes through the center point c. For more details on total equilibria and the motivating application, see [3].

Acknowledgment

We thank Raimund Seidel for suggesting the half-plane intersection approach which led to a considerable improvement over our original approach to output-sensitively exploring the zone of a curve in an arrangement of lines (Section 4) .

References

[1] K.-F. Böhringer. *Programmable Force Fields for Distributed Manipulation, and Their Implementation Using Microfabricated Actuator Arrays.* PhD thesis, Cornell University, Department of Computer Science, Ithaca, NY 14853, Aug. 1997.

[2] K.-F. Böhringer, V. Bhatt, and K. Y. Goldberg. Sensorless manipulation using transverse vibrations of a plate. In *Proc. IEEE Int. Conf. on Robotics and Automation (ICRA)*, pages 1989 – 1996, Nagoya, Japan, May 1995. `http://www.cs.cornell.edu/home/karl/VibratoryAlign`.

[4] Small disturbances from a *stable* equilibrium cause the part to return to this equilibrium. Slight disturbances in an *unstable* equilibrium are sufficient to move the part further away from equilibrium.

[3] K.-F. Böhringer, B. R. Donald, and N. C. MacDonald. Upper and lower bounds for programmable vector fields with applications to MEMS and vibratory plate parts feeders. In J.-P. Laumond and M. Overmars, editors, *Algorithms for Robotic Motion and Manipulation*, pages 255–276. A. K. Peters, Wellesley, MA 02181, 1997. `http://www.cs.cornell.edu/home/karl/MicroManipulation`.

[4] K.-F. Böhringer, B. R. Donald, N. C. MacDonald, G. T. A. Kovacs, and J. W. Suh. Computational methods for design and control of MEMS micromanipulator arrays. *Computer Science and Engineering*, pages 17 – 29, January – March 1997.

[5] T. K. Dey. Improved bounds for k-sets and k-th levels. In *Proc. IEEE Symp. on Foundations of Computer Science (FOCS)*. 1997. To appear.

[6] M. Díaz and J. O'Rourke. Ham-sandwich sectioning of polygons. In *Proc. 2nd Canadian Conference on Computational Geometry*, pages 98–101, Ottawa, 1990.

[7] M. G. Díaz. *Algorithms for Balanced Partitioning of Polygons and Point Sets*. PhD thesis, Johns Hopkins University, Baltimore, Maryland, 1990.

[8] H. Edelsbrunner. *Algorithms in Combinatorial Geometry*, volume 10 of *EATCS Monographs on Theoretical Computer Science*. Springer Verlag, Heidelberg, Germany, 1987.

[9] H. Edelsbrunner, L. Guibas, J. Pach, R. Pollack, R. Seidel, and M. Sharir. Arrangements of curves in the plane: Topology, combinatorics, and algorithms. *Theoretical Compututer Science*, 92:319–336, 1992.

[10] H. Edelsbrunner and E. Welzl. On the number of line separations of a finite set in the plane. *J. Combin. Theory Ser. A*, 1985.

[11] L. J. Guibas and J. Stolfi. Primitives for the manipulation of general subdivisions and the computation of Voronoi diagrams. *ACM Trans. Graph.*, 4, 1985.

[12] D. Halperin. Arrangements. In J. E. Goodman and J. O'Rourke, editors, *Handbook of Discrete and Computational Geometry*, chapter 21, pages 389–412. CRC Press LLC, 1997.

[13] S. Har-Peled. The complexity of many cells in the overlay of many arrangements. M.S. thesis, Dept. Computer Science, Tel Aviv University, Tel Aviv, Israel, 1995.

[14] J. O'Rourke. *Computational Geometry in C*. Cambridge University Press, 1994.

[15] M. H. Overmars and J. van Leeuwen. Maintenance of configurations in the plane. *J. Comput. Syst. Sci.*, 23:166–204, 1981.

[16] T. C. Shermer. A linear algorithm for bisecting a polygon. *Inform. Process. Lett.*, 41:135–140, 1992.

APPENDIX

The results presented in this paper make use of several geometric properties of polygon bisectors, as summarized in Section 2. In this appendix we present a complete derivation of all the formulas necessary to prove Propositions 2 and 3.

Consider a line λ, and a point s that lies on λ (Figure 2). The direction of λ is given by a vector p. Assume for now that the line λ intersects two edges e_1 and e_2 of the polygon P in the points p_1 and p_2. Also assume that these edges have direction vectors q_1 and q_2.

We will use the following notation throughout the appendix. For two vectors $v_1 = (x_1, y_1)$ and $v_2 = (x_2, y_2)$, $v_1 \times v_2$ will denote the signed real whose value is $x_1 y_2 - y_1 x_2$.

Rotating the Bisector. Consider another line λ' with direction p' that intersects λ in s (λ' can be understood as a rotation of λ around s). Assume that λ and λ' have combinatorially equivalent intersections with polygon P, and that λ' intersects the polygon edges e_1 and e_2 in p'_1 and p'_2 respectively. Let us write $p_i = s + \rho_i p$ and $p'_i = s + \rho'_i p'$. Then the polygon area a between λ and λ' is:

$$\begin{aligned} a &= \frac{1}{2}(\rho'_2 p' \times \rho_2 p - \rho'_1 p' \times \rho_1 p) \\ &= \frac{1}{2}(\rho'_2 \rho_2 - \rho'_1 \rho_1)(p' \times p) \end{aligned}$$

In the general case where λ and λ' intersect multiple edges of some arbitrary polygon P at points $p_1, p_2, \ldots, p_r$ and $p'_1, p'_2, \ldots, p'_r$ (r even), we can compute the polygon area a between λ and λ' as a sum of signed areas of triangles $sp_ip'_i$:

$$\begin{aligned} a &= \frac{1}{2}\sum_{i=1}^{r}(-1)^i w_i \\ &= \frac{1}{2}(p' \times p)\sum_{i=1}^{r}(-1)^i \rho'_i \rho_i \end{aligned}$$

Without loss of generality assume $\rho_r \neq 0$. Then p' can be written as $p' = p + \alpha q_r$ for some $\alpha \in \mathbb{R}$, and the above equation becomes:

$$\begin{aligned} a &= \frac{1}{2}((p + \alpha q_r) \times p)\sum_{i=1}^{r}(-1)^i \rho'_i \rho_i \\ &= \frac{\alpha}{2}(q_r \times p)\sum_{i=1}^{r}(-1)^i \rho'_i \rho_i \end{aligned} \tag{1}$$

From the two vector equations $p'_i = s + \rho'_i p'$ and $p'_i = s + \rho_i p + \mu q_i$, $\mu \in \mathbb{R}$, we can determine ρ'_i as:

$$\rho'_i = \frac{\rho_i (q_i \times p)}{(q_i \times p) + \alpha(q_i \times q_r)} \tag{2}$$

Note that the denominator in Eq. (2) can be written as $q_i \times (p + \alpha q_r) = q_i \times p' \neq 0$ since q_i cannot be parallel to p' as long as the intersections p_i lie in the same equivalence class. If we also choose the edge direction vectors q_i such that $(q_i \times p) = 1$, then Equations 2 and 1 simplify respectively to the following rational functions in α:

$$\rho'_i = \frac{\rho_i}{1 + \alpha(q_i \times q_r)} \tag{3}$$

$$a(\alpha) = \frac{\alpha}{2}\sum_{i=1}^{r}(-1)^i \frac{\rho_i^2}{1 + \alpha(q_i \times q_r)} \tag{4}$$

Let us look at the denominator $d_i(\alpha) = 1 + \alpha(q_i \times q_r)$ in more detail. This is important because we shall see that in the formulas that we will obtain, the denominators consist only of $d_i(\alpha)$'s. For an arbitrary polygon, d_i is a linear function of α. If all q_i are parallel then $d_i = 1$.

Equation 4 can be used to find a bisector of P that passes through a given point s. Given q_i and ρ_i ($i = 1 \ldots r$), a is chosen such that $|P \cap h_l(\lambda')| = |P \cap h_r(\lambda')| = \frac{1}{2}|P|$. Then Equation 4 can be transformed into a polynomial in α of degree r. More specifically, the degree of the polynomial α is equal to the number t of distinct slopes of polygon edges that the bisector crosses, which in the worst case is r, and c for a c-oriented polygon. This result was summarized in Proposition 2.1.

Translating the Bisector. We now consider the case where λ' shifts parallel (Figure 3). Analogously to the rotation case, let $p'_i = s' + \rho'_i p'$, and $p''_i = s'' + \rho''_i p'$. Also s' and s'' are chosen such that $s'' - s' = \alpha' q_2$. Then the polygon area between λ' and λ'' is:

$$\begin{aligned} b &= \alpha' q_2 \times \frac{1}{2}((p'_2 + p''_2) - (p'_1 + p''_1)) \\ &= \frac{\alpha'}{2}(\rho'_2 + \rho''_2 - \rho'_1 - \rho''_1)(q_2 \times (p + \alpha q_2)) \\ &= \frac{\alpha'}{2}(\rho'_2 + \rho''_2 - \rho'_1 - \rho''_1) \end{aligned} \tag{5}$$

In the general case λ' and λ'' intersect multiple edges of some arbitrary polygon P at points $p'_1, p'_2, \ldots, p'_r$ and $p''_1, p''_2, \ldots, p''_r$. We choose s' and s'' such that $s'' - s' = \alpha' q_r$. Now the ρ''_i can be determined from the two vector equations $p''_i = p'_i + \mu q_i$, $\mu \in \mathbb{R}$, and $p''_i = s'' + \rho''_i p'$:

$$\begin{aligned} \rho''_i &= \rho'_i - \alpha' \frac{q_i \times q_r}{q_i \times p'} \\ &= \rho'_i - \alpha' \frac{q_i \times q_r}{1 + \alpha(q_i \times q_r)} \\ &= \frac{\rho_i - \alpha'(q_i \times q_r)}{1 + \alpha(q_i \times q_r)} \end{aligned} \tag{6}$$

Then the polygon area between λ' and λ'' is:

$$\begin{aligned} b &= \frac{\alpha'}{2} \sum_{i=1}^{r} (-1)^i (\rho'_i + \rho''_i) \\ &= \frac{\alpha'}{2} \sum_{i=1}^{r} (-1)^i \frac{2\rho_i - \alpha'(q_i \times q_r)}{1 + \alpha(q_i \times q_r)} \end{aligned} \tag{7}$$

This is a quadratic polynomial in α' (unless all q_i are parallel, in which case it simplifies to the linear equation $b = \alpha' \sum_{i=1}^{r} (-1)^i \rho_i$). Equation 7 can be used to find a bisector of P with a given direction p, i.e., with a given slope. This result was summarized in Proposition 2.2.

Maintaining the Bisector Property. From the analysis so far we see that if the bisector λ is rotated to λ', then the left and right areas are changed by a value a ($\neq 0$ in general) as described in Equation 4. Hence a subsequent shift of λ' is necessary to restore the bisector property, by changing the areas by a value b, as described in Equation 7.

This implies the condition $a + b = 0$, with a and b given by Equations 4 and 7:

$$\begin{aligned} a + b &= \frac{1}{2} \sum_{i=1}^{r} (-1)^i \frac{\alpha \rho_i^2 + 2\alpha' \rho_i - (\alpha')^2 (q_i \times q_r)}{1 + \alpha(q_i \times q_r)} \\ &= 0 \end{aligned} \tag{8}$$

This equation ensures that λ is a bisector of P. Equation 8 describes $a + b$ as a rational function in α, and a quadratic polynomial in α'. Hence for all combinatorially equivalent bisectors, we can obtain an explicit formula to describe α' as a function of α. We conclude that all combinatorially equivalent bisectors can be represented by an explicit expression parameterized by α.

In general, Equation 8 is equivalent to a polynomial in α and α' whose degree depends on the number $t \leq r$ of distinct slopes of polygon edges intersected by the bisectors λ, λ', or λ'', provided that the $d_i(\alpha)$'s are all non-zero; see remark above. The polynomial is at most quadratic in α'. In the case that all q_i are parallel, Equation 8 simplifies to the linear equation $\sum_{i=1}^{r} (-1)^i \left(\alpha \frac{\rho_i}{2} + \alpha'\right) \rho_i = 0$. These results were summarized in Proposition 3.