

Section 3.5

Which of the following languages are CFL?

- $L = \{a^n b^n c^j \mid 0 < n \leq j\}$
- $L = \{a^n b^j a^n b^j \mid n > 0, j > 0\}$
- $L = \{a^n b^j a^k b^p \mid n + j \leq k + p, n > 0, j > 0, k > 0, p > 0\}$

Pumping Lemma for Regular Language's: Let L be a regular language, Then there is a constant m such that $w \in L$, $|w| \geq m$, $w = xyz$ such that

- $|xy| \leq m$
- $|y| \geq 1$
- for all $i \geq 0$, $xy^i z \in L$

Pumping Lemma for CFL's Let L be any infinite CFL. Then there is a constant m depending only on L , such that for every string w in L , with $|w| \geq m$, we may partition $w = uvxyz$ such that:

$$\begin{aligned} |vxy| &\leq m, \text{ (limit on size of substring)} \\ |vy| &\geq 1, \text{ (} v \text{ and } y \text{ not both empty)} \\ \text{For all } i &\geq 0, uv^i xy^i z \in L \end{aligned}$$

- **Proof:** (sketch) There is a CFG G s.t. $L=L(G)$.

Consider the parse tree of a long string in L .

For any long string, some nonterminal N must appear twice in the path.

Example: Consider $L = \{a^n b^n c^n : n \geq 1\}$. Show L is not a CFL.

• **Proof:** (by contradiction)

Assume L is a CFL and apply the pumping lemma.

Let m be the constant in the pumping lemma and consider $w = a^m b^m c^m$. Note $|w| \geq m$.

Show there is no division of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$, and $uv^i xy^i z \in L$ for $i = 0, 1, 2, \dots$

Case 1: Neither v nor y can contain 2 or more distinct symbols. If v contains a 's and b 's, then $uv^2 xy^2 z \notin L$ since there will be b 's before a 's.

Thus, v and y can be only a 's, b 's, or c 's (not mixed).

Case 2: $v = a^{t_1}$, then $y = a^{t_2}$ or b^{t_3} ($|vxy| \leq m$)

If $y = a^{t_2}$, then $uv^2 xy^2 z = a^{m+t_1+t_2} b^m c^m \notin L$ since $t_1 + t_2 > 0$, $n(a) > n(b)$'s (number of a 's is greater than number of b 's)

If $y = b^{t_3}$, then $uv^2 xy^2 z = a^{m+t_1} b^{m+t_3} c^m \notin L$ since $t_1 + t_3 > 0$, either $n(a) > n(c)$'s or $n(b) > n(c)$'s.

Case 3: $v = b^{t_1}$, then $y = b^{t_2}$ or c^{t_3}

If $y = b^{t_2}$, then $uv^2 xy^2 z = a^m b^{m+t_1+t_2} c^m \notin L$ since $t_1 + t_2 > 0$, $n(b) > n(a)$'s.

If $y = c^{t_3}$, then $uv^2 xy^2 z = a^m b^{m+t_1} c^{m+t_3} \notin L$ since $t_1 + t_3 > 0$, either $n(b) > n(a)$'s or $n(c) > n(a)$'s.

Case 4: $v = c^{t_1}$, then $y = c^{t_2}$

then, $uv^2 xy^2 z = a^m b^m c^{m+t_1+t_2} \notin L$ since $t_1 + t_2 > 0$, $n(c) > n(a)$'s.

Thus, there is no breakdown of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$ and for all $i \geq 0$, $uv^i xy^i z$ is in L . Contradiction, thus, L is not a CFL. Q.E.D.

Example Why would we want to recognize a language of the type $\{a^n b^n c^n : n \geq 1\}$?

Example: Consider $L = \{a^n b^n c^p : p > n > 0\}$. Show L is not a CFL.

- **Proof:** Assume L is a CFL and apply the pumping lemma. Let m be the constant in the pumping lemma and consider $w = \underline{\hspace{2cm}}$. Note $|w| \geq m$.
Show there is no division of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$, and $uv^i xy^i z \in L$ for $i = 0, 1, 2, \dots$

Thus, there is no breakdown of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$ and for all $i \geq 0$, $uv^i xy^i z$ is in L . Contradiction, thus, L is not a CFL. Q.E.D.

Example: Consider $L = \{a^j b^k : k = j^2\}$. Show L is not a CFL.

- **Proof:** Assume L is a CFL and apply the pumping lemma. Let m be the constant in the pumping lemma and consider $w = \underline{\hspace{2cm}}$

Show there is no division of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$, and $uv^i xy^i z \in L$ for $i = 0, 1, 2, \dots$

Case 1: Neither v nor y can contain 2 or more distinct symbols. If v contains a 's and b 's, then $uv^2 xy^2 z \notin L$ since there will be b 's before a 's.

Thus, v and y can be only a 's, and b 's (not mixed).

Thus, there is no breakdown of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$ and for all $i \geq 0$, $uv^i xy^i z$ is in L . Contradiction, thus, L is not a CFL. Q.E.D.

Exercise: Prove the following is not a CFL by applying the pumping lemma. (answer is at the end of this handout).

Consider $L = \{a^{2^n} b^{2^p} c^n d^p : n, p \geq 0\}$. Show L is not a CFL.

Example: Consider $L = \{w\bar{w}w : w \in \Sigma^*\}$, $\Sigma = \{a, b\}$, where \bar{w} is the string w with each occurrence of a replaced by b and each occurrence of b replaced by a . For example, $w = baaa$, $\bar{w} = abbb$, $w\bar{w} = baaaabbb$. Show L is not a CFL.

- **Proof:** Assume L is a CFL and apply the pumping lemma. Let m be the constant in the pumping lemma and consider $w = \underline{\hspace{2cm}}$

Show there is no division of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$, and $uv^i xy^i z \in L$ for $i = 0, 1, 2, \dots$

Thus, there is no breakdown of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$ and for all $i \geq 0$, $uv^i xy^i z$ is in L . Contradiction, thus, L is not a CFL. Q.E.D.

Example: Consider $L = \{a^n b^p b^p a^n\}$. L is a CFL. The pumping lemma should apply!

Let $m \geq 4$ be the constant in the pumping lemma. Consider $w = a^m b^m b^m a^m$.

We can break w into $uvxyz$, with:

If you apply the pumping lemma to a CFL, then you should find a partition of w that works!

Chap 8.2 Closure Properties of CFL's

Theorem CFL's are closed under union, concatenation, and star-closure.

- **Proof:**

Given 2 CFG $G_1 = (V_1, T_1, R_1, S_1)$ and $G_2 = (V_2, T_2, P_2, S_2)$

- Union:

Construct G_3 s.t. $L(G_3) = L(G_1) \cup L(G_2)$.

$G_3 = (V_3, T_3, R_3, S_3)$

- Concatenation:

Construct G_3 s.t. $L(G_3) = L(G_1) \circ L(G_2)$.

$G_3 = (V_3, T_3, R_3, S_3)$

- Star-Closure
Construct G_3 s.t. $L(G_3) = L(G_1)^*$
 $G_3 = (V_3, T_3, R_3, S_3)$

QED.

Theorem CFL's are NOT closed under intersection and complementation.

- **Proof:**

- Intersection:

- Complementation:

Theorem: CFL's are closed under *regular* intersection. If L_1 is CFL and L_2 is regular, then $L_1 \cap L_2$ is CFL.

- **Proof:** (sketch) This proof is similar to the construction proof in which we showed regular languages are closed under intersection. We take a NPDA for L_1 and a DFA for L_2 and construct a NPDA for $L_1 \cap L_2$.

$M_1 = (Q_1, \Sigma, \delta_1, q_0, z, F_1)$ is an NPDA such that $L(M_1) = L_1$.

$M_2 = (Q_2, \Sigma, \delta_2, q'_0, F_2)$ is a DFA such that $L(M_2) = L_2$.

Example of replacing arcs (NOT a Proof!):

Note this is not a proof, but sketches how we will combine the DFA and NPDA. We must formally define Δ_3 . If

then

Must show

if and only if

Must show:

$w \in L(M_3)$ iff $w \in L(M_1)$ and $w \in L(M_2)$.

QED.

Questions about CFL:

1. Decide if CFL is empty?

2. Decide if CFL is infinite?

Example: Consider $L = \{a^{2n}b^{2m}c^nd^m : n, m \geq 0\}$. Show L is not a CFL.

- **Proof:** Assume L is a CFL and apply the pumping lemma. Let m be the constant in the pumping lemma and consider $w = a^{2m}b^{2m}c^md^m$.

Show there is no division of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$, and $uv^ixy^iz \in L$ for $i = 0, 1, 2, \dots$

Case 1: Neither v nor y can contain 2 or more distinct symbols. If v contains a 's and b 's, then $uv^2xy^2z \notin L$ since there will be b 's before a 's.

Thus, v and y can be only a 's, b 's, c 's, or d 's (not mixed).

Case 2: $v = a^{t_1}$, then $y = a^{t_2}$ or b^{t_3} ($|vxy| \leq m$)

If $y = a^{t_2}$, then $uv^2xy^2z = a^{2m+t_1+t_2}b^{2m}c^md^m \notin L$ since $t_1 + t_2 > 0$, the number of a 's is not twice the number of c 's.

If $y = b^{t_3}$, then $uv^2xy^2z = a^{2m+t_1}b^{2m+t_3}c^md^m \notin L$ since $t_1 + t_3 > 0$, either the number of a 's (denoted $n(a)$) is not twice $n(c)$ or $n(b)$ is not twice $n(d)$.

Case 3: $v = b^{t_1}$, then $y = b^{t_2}$ or c^{t_3}

If $y = b^{t_2}$, then $uv^2xy^2z = a^{2m}b^{2m+t_1+t_2}c^md^m \notin L$ since $t_1 + t_2 > 0$, $n(b) > 2*n(d)$.

If $y = c^{t_3}$, then $uv^2xy^2z = a^{2m}b^{2m+t_1}c^{m+t_3}d^m \notin L$ since $t_1 + t_3 > 0$, either $n(b) > 2*n(d)$ or $2*n(c) > n(a)$.

Case 4: $v = c^{t_1}$, then $y = c^{t_2}$ or d^{t_3}

If $y = c^{t_2}$, then $uv^2xy^2z = a^{2m}b^{2m}c^{m+t_1+t_2}d^m \notin L$ since $t_1 + t_2 > 0$, $2*n(c) > n(a)$.

If $y = d^{t_3}$, then $uv^2xy^2z = a^{2m}b^{2m}c^{m+t_1}d^{m+t_3} \notin L$ since $t_1 + t_3 > 0$, either $2*n(c) > n(a)$ or $2*n(d) > n(b)$.

Case 5: $v = d^{t_1}$, then $y = d^{t_2}$

then $uv^2xy^2z = a^{2m}b^{2m}c^md^{m+t_1+t_2} \notin L$ since $t_1 + t_2 > 0$, $2*n(d) > n(c)$.

Thus, there is no breakdown of w into $uvxyz$ such that $|vy| \geq 1$, $|vxy| \leq m$ and for all $i \geq 0$, uv^ixy^iz is in L . Contradiction, thus, L is not a CFL. Q.E.D.